

Palbociclib (in combination with fulvestrant)

Indication

Treatment of hormone receptor-positive, human epidermal growth factor receptor 2 (HER2)-negative, locally advanced or metastatic breast cancer in people who have had previous endocrine therapy only if exemestane plus everolimus is the most appropriate alternative to a cyclin- dependent kinase 4 and 6 (CDK 4/6) inhibitor.

(NICE TA619)

ICD-10 codes

Codes with a pre fix C50.

Regimen details

Day	Drug	Dose	Route
1-21	Palbociclib	125mg OD	PO

Fulvestrant 500 mg should be administered intramuscularly on days 1 and 15 of cycle 1 and on day 1 of subsequent cycles.

Cycle frequency

28 days.

Palbociclib should be taken for 21 days followed by a 7 day break.

Number of cycles

Until disease progression or unacceptable toxicity.

Administration

Palbociclib is available as 125mg, 100mg and 75mg capsules. The capsules should be swallowed whole and not chewed, crushed or opened. The dose should be taken with food, preferably a meal.

Grapefruit and grapefruit juice should be **avoided** whilst taking palbociclib.

Patients should be advised to take the dose at approximately the same time each day. If a patient vomits or misses a dose an additional dose should not be taken that day but the next prescribed dose should be taken as planned.

Pre-medication

Nil

Emetogenicity

This regimen has mild emetic potential.

Additional supportive medication

Nil

Extravasation

N/A

Investigations – pre first cycle

Investigation	Validity period
FBC	14 days
U+Es (including creatinine)	14 days
LFTs	14 days

Investigations – pre subsequent cycles

Investigation	Validity period
FBC	72 hours and on day 14 for the first 2 cycles*
U+Es (including creatinine)	72 hours
LFTs	72 hours

*If neutrophils $< 1.0 \times 10^9/L$ or platelets $< 50 \times 10^9/L$ where possible repeat on day 1 of planned cycle.

Standard limits for administration to go ahead

If blood results not within range, authorisation to administer **must** be given by prescriber/ consultant

Investigation	Limit
Neutrophils	$\geq 1.0 \times 10^9/L$
Platelets	$\geq 50 \times 10^9/L$
CrCl	$\geq 30\text{mL}/\text{min}$
Bilirubin	$< 1.5 \times \text{ULN}$
AST/ALT	$< \text{ULN}$

Dose modifications

Dose reductions should follow the table below:

Dose level	Dose
Full dose	125mg OD
First reduction	100mg OD
Second reduction	75mg OD

Dose reductions below 75mg OD are not recommended and if required treatment should be discontinued.

- **Haematological toxicity**

On day 1 neutrophils $\geq 1.0 \times 10^9/L$ and platelets $\geq 50 \times 10^9/L$.

Dose interruption, dose reduction, or delay in starting treatment cycles is recommended for patients who develop grade 3 or 4 neutropenia.

Haematological toxicity	Dose
Grade 1-2 (neutrophils $\geq 1.0 \times 10^9/L$)	No dose modification required.
Grade 3 (neutrophils $0.5- 1.0 \times 10^9/L$)	Day 1: Withhold and repeat FBC. When recovered to \leq grade 2 start next cycle at the same dose). Day 14 (cycles 1 and 2): Continue to complete cycle and repeat FBC on day 21. Consider dose reduction if not recovered within 7 days or recurrent neutropenia.
Grade 3 (neutrophils $0.5- 1.0 \times 10^9/L$) with fever +/- infection	Withhold until recovered to \leq grade 2. Resume with one dose level reduction.
Grade 4 (neutrophils $< 0.5 \times 10^9/L$)	Withhold until recovered to \leq grade 2. Resume with one dose level reduction.

- **Renal impairment**

Palbociclib should be administered with caution and close monitoring for signs of toxicity in severe renal impairment (CrCl <30mL/min).

- **Hepatic impairment**

Palbociclib should be administered with caution in moderate to severe hepatic impairment (bilirubin > 1.5 x ULN and/or AST/ALT > ULN). The risk and benefits should be carefully considered and patients should be closely monitored for signs of toxicity.

- **Other toxicities**

For any other non-haematological toxicity ≥ Grade 3; withhold until ≤ Grade 1 (≤ Grade 2 if not considered safety risk) then resume with one dose level reduction.

Adverse effects - for full details consult product literature/ reference texts

- **Serious side effects**

Neutropenia, anaemia, leukopenia.
Infections

- **Frequently occurring side effects**

Neutropenia, anaemia, leukopenia.
Thrombocytopenia
Infections
Fatigue
Nausea and vomiting
Stomatitis
Rash, dry skin
Alopecia
Diarrhoea

- **Other side effects**

Reduced appetite
Dysgeusia
Blurred vision
Dry eyes
Increased transaminases

Significant drug interactions – for full details consult product literature/ reference texts

Strong CYP3A4 inhibitors (e.g. clarithromycin, indinavir, itraconazole, ketoconazole, lopinavir/ritonavir, nefazodone, nelfinavir, posaconazole, saquinavir, telaprevir, telithromycin, voriconazole, grapefruit): Concomitant use of strong inhibitors should be avoided due to increased risk of toxicity. If co-administrated is deemed essential the dose of palbociclib should be reduced to 75mg daily and patients closely monitored.

Strong CYP3A4 inducers (e.g. carbamazepine, enzalutamide, phenytoin, rifampin, and St. John's Wort): Concomitant use may reduce the exposure of palbociclib and should therefore be avoided.

Additional comments

Women of childbearing potential or their male partners must use a highly effective method of contraception.

This medicinal product contains lactose. Patients with rare hereditary problems of galactose intolerance, the Lapp lactase deficiency, or glucose-galactose malabsorption should not take this medicine.

References

- National Institute for Clinical Excellence (TA619) accessed January 2020 via www.nice.org.uk
- Summary of Product Characteristics Palbociclib (Pfizer) accessed January 2020 via www.medicines.org.uk
- Turner M., et al. Overall Survival with Palbociclib and Fulvestrant in Advanced Breast Cancer. N Engl J Med 2018; 379:1926-1936

Written/reviewed by: Dr R Bowen (Consultant Oncologist, RUH Bath NHS Trust), Dr M Beresford (Consultant Oncologist, RUH Bath NHS Trust)

Checked by: Sarah Murdoch (Oncology/Haematology Pharmacist, SW Clinical Network)

Authorised by: Dr J Braybrooke (Consultant Oncologist, SW Clinical Network)

Date: January 2020
